

An Odyssey of discovery: principles of agriculture for the humid tropics

Roland Bunch

Our Odyssey started in 1982, the day Conrado, a Honduran villager, sheepishly showed us his experiment. Sceptical about the value of the organic matter we had recommended, he had piled a huge quantity of compost into several rows of his maize field. The last two rows he left as a control, untilled and unfertilized. There, before our eyes, stood a field of maize 2.5 meters tall, with a last row reaching not even 40 cms. That was the day we began to realize the incredible degree to which organic matter can restore degraded soils.

Little by little, work in a dozen countries has convinced us that the vast majority of soils can be made highly fertile. How? By using our first principle: maximize organic matter production. Conrado's particular approach, however, was anti-economic. The cost of using compost on basic grains exceeds the benefit. But intercropped green manure/cover crops (gm/cc's) can produce from 50 to 140 t/ha (green weight) of organic matter with very little work: no transporting of material and no cutting up or layering or turning over of compost heaps. In fact, sometimes, because of the gm/cc's control of weeds, net labour costs decrease, And soil quality often improves visibly each year.

Farmers find the answer

Then, as often happens, we found we were far from the first to employ intercropped gm/cc's. Gradually, between 1985 and 1992, we learned that villager farmers spread from Veracruz State in Mexico through Guatemala, El Salvador, and Honduras are intercropping velvetbeans (*Mucuna ensiformis*) with their maize and sorghum. To our amazement, these systems, virtually all of them in the supposedly infertile humid tropics, allow farmers to plant maize every year for decades, with productivity increasing over time up to 4 t/ha (Durón, 1990). In other words, these farmers have found an answer to slash-and-burn agriculture.

Migratory agriculture is most frequently motivated by decreasing fertility, increased weed problems, or both. In the Mesoamerican gm/cc systems, nitrogen fixation and biomass recycling maintain soil fertility. Mulches of crop residues and fast-growing gm/cc's drastically reduce the weed problem. We had learned a second principle: keep the soil covered.

Gm/cc mulches provide a whole series of additional benefits. They protect the soil from irradiation and the heat of the tropical sun, thereby also reducing organic matter burn-out. They save a tremendous amount of work; incorporating gm/cc's is virtually impossible for farmers using hand hoes. They keep the excess nitrogen from acidifying the upper soil horizons (Triomphe, 1994). And they largely prevent soil erosion, even on slopes of 40%.

Zero tillage

In the meantime, we had been reading Fukuoka's book, *The One Straw Revolution* (Fukuoka, 1978). However, his recommendation of zero tillage failed to convince us. After all, most of the traditional agriculture in Latin America uses zero tillage, yet is far from productive.

In mid-1993, I visited the work of EPAGRI in southern Brazil. Having visited over 160 agricultural development programmes through the years, I found this largely unpublicized effort to be the finest of its size I had seen in Latin America. Literally tens of thousands of animal traction farmers were producing harvests approaching those in the USA, with gm/cc's and zero tillage (Bunch, nd).

Valdemar de Freitas, EPAGRI's manager, showed us that the secret to achieving zero tillage is applying massive amounts of organic matter to the soil. Brazilian farmers, after some four years of applying gm/cc's to the soil, are able to quit ploughing. The advantages, in terms of better soil structure, reduced soil compaction, higher fertility and decreases cost, are impressive. Interestingly, farmers often use non-leguminous gm/cc's to increase biomass in order to quit ploughing sooner. That is, they spend scarce income on nitrogen for three or four years in order to achieve zero tillage sooner.

The Brazilians' discovery explains why the zero tillage gm/cc systems of northern Honduras, and Fukuoka's, produce so well, while many traditional zero tillage systems do not. Thus we added a third principle: use zero tillage.

Diversity

EPAGRI's investigation and dissemination of over 60 species of gm/cc's, partly to avoid diseases and insect pests, confirmed another, more widely known principle: maintain biological diversity.

The last principle was discovered by Martha Rosemeyer, a Cornell doctoral candidate working in Costa Rica. For several years, agronomists working with a low-cost traditional mulched bean (*Phaseolus vulgaris*) system had been trying to solve a phosphorus deficiency problem. With highly acid (pH = 4.0 to 4.5) soils, virtually all the phosphorus applied became tied up almost instantly. Farmers' harvests averaged 500 kgs/ha (Rosemeyer, 1994). Martha and a group of farmers tried broadcasting the phosphorus on top of the mulch.

The results, since confirmed in numerous additional experiments, were astounding. Bean yields rose to between 1.5 and 2.5 t/ha (Schlater, 1995). This phenomenon has not yet been validated with other crops. Yet it would help explain the success of Mesoamerica's gm/cc systems, and coincides with the fact that plants as diverse as maize, manioc and tropical trees tend to develop a heavy mass of feeder roots immediately under thick mulches (Rattan Lal, 1989; Vitousek and Sanford, 1986).

Furthermore, it makes simple sense: when soils are as hostile to plant growth as are the humid tropic's acid soils, feeding plants through a mulch would seem a much more promising alternative. The fifth principle is undoubtedly the most unconventional: feed plants through the mulch.

Imitating nature

These five principles enjoy a nice synergy. For example, if we are going to feed our plants through a mulch, we certainly cannot plough our fields. Nevertheless, the most important relation between these principles is precisely the one that took us the longest to figure out: they describe quite well the way a humid tropical forest functions.

That is, all we discovered in our 12 year Odyssey is something we should have guessed all along. In order for humid tropical agriculture to be both highly productive and sustainable, it must imitate the highly productive, millions-of-years-old humid tropical forest.

Three months ago, I searched the computerized agricultural data system in the United States for information on the nutrient dynamics in mulches and the feeding of crops through a mulch. I found virtually nothing. The above principles mean we are going to have to develop agricultural systems totally different from those agronomists have tried, for so many years, to "transfer" from the temperate nations.

Yet the possibilities are enormous. A study from northern Honduras shows that the gm/cc maize system there is 30% more profitable than the high-input maize system nearby (Milton Flores, 1992). It may well be we are just beginning to fathom the full potential of low-input agriculture in the humid tropics.

Roland Bunch, COSECHA, Apartado 3586, Tegucigalpa, Honduras.

References

- Durón, Elio. 1990. Presentation made to CIDICCO's First Interchange of Ideas on the Role of Leguminous Plans in Today's Agriculture, Tegucigalpa, Honduras.
- Triomphe, Bernard. 1994. Personal communication on the results of his doctoral dissertation research on the long-term effects of a Honduran gm/cc maize system on tropical soils.
- Fukuoka, Masanobu. 1978. *The one straw revolution: an introduction to natural farming*. Emmaus, Pennsylvania: Rodale Press.
- Bunch, Roland. nd. *EPAGRI's Work in the State of Santa Catarina, Brazil: Major new Possibilities for Resource-Poor Farmers*. photocopied.
- Rosemeyer, Martha. 1994. *Yield, nodulation and mycorrhizal establishment in slash/mulch vs. row-cropped beans*. In: H. David Thurston, et al. (eds). *Tapado, slash/mulch: how farmers use it and what researchers know about it*". pp. 169–178. Ithaca, NY: CIIFAD and CATIE.
- Schlater, Kenneth. 1995. Personal communication on the results of his doctoral dissertation research on the effects of applying phosphorus to the mulch of slash/mulched beans.
- Rattan Lal. 1989. *Conservation tillage for sustainable agriculture: tropics vs. temperate environments*. In: NC Brady (ed.). *Advances in Agronomy Vol. 42*. San Diego, California: Academic Press.
- PM Vitousek and RL Sanford, Jr. 1986. *Nutrient cycling in moist tropical forest*. In: *Annual Review of Ecological Systems, vol. 17, pp. 137–167*.
- Milton Flores and Nicolas Estrada. 1992. *Estudio de caso: la utilización del frijol abono (Mucuna spp.) como alternativa viable para el sostenimiento productivo de los sistemas agrícolas del litoral Atlántico*. Paper presented to the Center for Development Studies at the Free University of Amsterdam. Mimeographed.

[Back to Top](#)